

## IN THE CLAIMS

Please amend the claims as shown below.

1. (Currently Amended) A modified thermostable DNA polymerase ~~obtained~~ obtainable from *Pyrococcus furiosus*, ~~*Pyrococcus kodakaraensis* KOD1~~ or *Thermococcus litoralis* ~~genus~~ having a 3'-5' exonuclease activity, wherein the modification is the replacement of histidine (H) by another amino acid in the DIETLYH or DIETFYH ~~DX<sub>1</sub>EX<sub>2</sub>X<sub>3</sub>X<sub>4</sub>H~~ sequence (D: aspartic acid, I: isoleucine, E: glutamic acid, T: threonine, L: leucine, F: phenylalanine, Y: tyrosine, H: histidine, ~~X<sub>1</sub>, X<sub>2</sub>, X<sub>3</sub> and X<sub>4</sub>: any amino acid~~) within the exonuclease I region of the thermostable DNA polymerase.

2. (Currently Amended) The modified thermostable DNA polymerase according to claim 1, wherein in the DIETLYH or DIETFYH ~~DX<sub>1</sub>EX<sub>2</sub>X<sub>3</sub>X<sub>4</sub>H~~ sequence, histidine (H) has been replaced by an amino acid selected from the group consisting of aspartic acid, glutamic acid, tyrosine, alanine, lysine and arginine.

3. (Original) The modified thermostable DNA polymerase according to claim 1 having the following physicochemical properties:

- (1) DNA extension rate: at least 20 bases/second; and
- (2) thermostability: it is capable of retaining 10% or more DNA polymerase activity of untreated DNA polymerase at pH 8.8 (determined at 25°C) after treatment at 95°C for 6 hours.

4. (Currently Amended) A modified thermostable DNA polymerase having a 3'-5' exonuclease activity and the following physicochemical properties:

- (1) DNA extension rate: at least 30 bases/second;
- (2) thermostability: the modified thermostable DNA polymerase being capable of retaining 40% or more DNA polymerase activity of untreated DNA polymerase at pH 8.8 (determined at 25°C) after treatment at 95°C for 6 hours; and
- (3) amino acid sequence: the modified thermostable DNA polymerase comprising an amino acid sequence of SEQ ID NO:2 except that ~~in the DX<sub>1</sub>EX<sub>2</sub>X<sub>3</sub>X<sub>4</sub>H amino acid sequence located at the 141- to 146-positions of SEQ ID NO:2 is selected from DIETLY or DIETFY and 147-positions in the amino acid sequence of SEQ ID NO: 2, histidine (H) at the 147-position of SEQ ID NO:2 has been replaced by another amino acid, wherein X<sub>1</sub> is isoleucine, X<sub>2</sub> is threonine, X<sub>3</sub> is leucine and X<sub>4</sub> is tyrosine.~~

5. (Previously Presented) The modified thermostable DNA polymerase according to claim 4 having the following thermostability: it is capable of retaining 60% or more DNA polymerase activity of untreated DNA polymerase at pH 8.8 (determined at 25°C) after treatment at 95°C for 6 hours.

6. (Currently Amended) The modified thermostable DNA polymerase according to claim 5, wherein ~~in the amino acid sequence of SEQ ID NO: 2~~, histidine (H) at the 147-position has been replaced by an amino acid selected from the group consisting of aspartic acid, glutamic acid, tyrosine, alanine, lysine and arginine.

7. (Currently Amended) The modified thermostable DNA polymerase according to claim 6, wherein ~~in the amino acid sequence of SEQ ID NO: 2~~, histidine (H) at the 147-position has been replaced by aspartic acid.

8. (Currently Amended) The modified thermostable DNA polymerase according to claim 6, wherein ~~in the amino acid sequence of SEQ ID NO: 2~~, histidine (H) at the 147-position has been replaced by glutamic acid.

9. (Currently Amended) The modified thermostable DNA polymerase according to claim 6, wherein ~~in the amino acid sequence of SEQ ID NO: 2~~, histidine (H) at the 147-position has been replaced by tyrosine.

10. (Currently Amended) The modified thermostable DNA polymerase according to claim 6, wherein ~~in the amino acid sequence of SEQ ID NO: 2~~, histidine (H) at the 147-position has been replaced by alanine.

11. (Currently Amended) The modified thermostable DNA polymerase according to claim 6, wherein ~~in the amino acid sequence of SEQ ID NO: 2~~, histidine (H) at the 147-position has been replaced by lysine.

12. (Currently Amended) The modified thermostable DNA polymerase according to claim 6, wherein ~~in the amino acid sequence of SEQ ID NO: 2~~, histidine (H) at the 147-position has been replaced by arginine.

13-24. (Canceled)

25. (Previously Presented) A reagent kit for amplifying nucleic acid, which comprises 2 kinds of primers, each of the primers being complementary to a DNA extension product of the other primer; dNTP; the thermostable DNA polymerase of claim 1; divalent ion(s); monovalent ion(s); and a buffer solution.

26. (Previously Presented) A reagent kit for amplifying nucleic acid, which comprises 2 kinds of primers, each of the primers being complementary to a DNA extension product of the other primer; dNTP; the thermostable DNA polymerase of claim 1; magnesium ion; at least one of monovalent ions selected from the group consisting of ammonium ion and potassium ion; BSA (bovine serum albumin); a nonionic surfactant and a buffer solution.

27. (Previously Presented) A reagent kit for amplifying nucleic acid, which comprises 2 kinds of primers, each of the primers being complementary to a DNA extension product of the other primer; dNTP; the thermostable DNA polymerase of claim 1; magnesium ion; at least one of monovalent ions selected from the group consisting of ammonium ion and potassium ion; BSA (bovine serum albumin); a nonionic surfactant; a buffer solution and an antibody capable of suppressing at least one activity selected from polymerase activity and 3'-5' exonuclease activity of the thermostable DNA polymerase.

28. (Previously Presented) A DNA polymerase composition which comprises one or more kinds of modified thermostable DNA polymerases defined in claim 1.

29. (Canceled)

30. (Previously Presented) A reagent kit for producing a mutated DNA which comprises mutagenesis primers, dNTP and the thermostable DNA polymerase of claim 1.

31. (Canceled)

32. (Previously Presented) A modified thermostable DNA polymerase according to claim 1 wherein said DNA polymerase is an  $\alpha$ -like DNA polymerase.

33. (Previously Presented) A modified thermostable DNA polymerase according to claim 1 wherein said DNA polymerase is thermostable DNA polymerase.

34-35. (Canceled)

36. (Previously Presented) A modified thermostable DNA polymerase according to claim 1 wherein the histidine (H) has been replaced by an acidic amino acid to obtain the modified thermostable DNA polymerase having significantly reduced 3'-5' exonuclease activity as compared with the enzyme before modification.

37. (Previously Presented) A modified thermostable DNA polymerase according to claim 1 wherein the histidine (H) has been replaced by a neutral amino acid to obtain a modified thermostable DNA polymerase having improved amplifying efficiency.

38. (Previously Presented) A modified thermostable DNA polymerase according to claim 1 wherein the histidine (H) has been replaced by a basic amino acid to obtain a modified thermostable DNA polymerase having significantly improved 3'-5' exonuclease activity and/or fidelity on a DNA replication or amplification.

39. (Previously Presented) The modified thermostable DNA polymerase according to claim 1, wherein the histidine (H) has been replaced by an acidic amino acid to obtain the modified thermostable DNA polymerase having improved PCR amplification efficiency from low copy number of template DNA.

40. (Previously Presented) The modified thermostable DNA polymerase according to claim 1, wherein the histidine (H) has been replaced by an acidic amino acid to obtain the modified thermostable DNA polymerase having improved PCR amplification efficiency from a long DNA segment.

41. (Previously Presented) The modified thermostable DNA polymerase according to claim 1, wherein the histidine (H) has been replaced by a neutral amino acid to obtain the modified thermostable DNA polymerase having improved PCR amplification efficiency from low copy number of template DNA.